

The Eldor Disposable Infusion Pump

Directions for Use



ISO 9001 & ISO 13485

Indications for Use

- The Eldor pump is indicated for delivery of medications
 - o Percutaneous Around surgical sites
 - o Intravenously
 - Via Central vascular access
- The Eldor pump is supplied sterile non pyrogenic
- The Eldor pump is for single use only
- Store out of direct sunlight

Cautions

- Do not use if package is damaged or open
- Do not resterilize or exceed maximum fill volume of pump
- The Eldor pump can exhibit a variance in accuracy of approximately + or 10 %
- Nominal infusion rate is labeled on the inside of each device
- Infusion time can be influenced by the viscosity of medication concentration
- Use strict aseptic technique when filling device

ALLIED MEDICAL LIMITED ABN: 99 089 052 479
LEVEL 2, 87 ADELAIDE TERRACE, EAST PERTH 6004
PO BOX 6915, EAST PERTH 6892
FAX: +61 8 9226 1999 info@

PH: +61 8 9266 0190

info@alliedmedical.com.au

Filling the Eldor Infusion pump

- 1. Remove device from packaging
- 2. Remove white cap (A) from filling port on top of device and keep
- 3. Slip the one way filter that is supplied securely into refill port
- **4.** Inject the desired volume of fluid via a syringe into device via one way filter. The elastomeric balloon will expand as solution is injected
- 5. Stand device will refill port at top, remove one way filter and expel any air that may have entered the balloon
- **6.** When balloon is full as per the required volume and free of air, securely replace white cap on top of refill port
- **7. To Prime the patient line -** remove cap from end of patient line and wait unit first drop of solution appears
- 8. Attach medication label and connect to patient

